

DEFY LINES.

(PARENTHESES HAVE NO PLACE)
ON YOUR FACE.



Actual patient. Results may vary.

- **Instantly** smooths away the deeper lines along the sides of your nose and mouth
- Provides **natural-looking** results
- **Lasts up to 1 year** with optimal treatment

APPROVED USE

JUVÉDERM® XC injectable gel is for injection into areas of facial tissue where moderate to severe facial wrinkles and folds occur to temporarily add volume to the skin, especially around the nose and mouth.

IMPORTANT SAFETY INFORMATION

Are there any reasons why I should not receive JUVÉDERM® XC?

Do not use the product if you have severe allergies with a history of severe allergic reactions (anaphylaxis), or if you are allergic to lidocaine or the proteins (gram-positive bacterial proteins) used to make the hyaluronic acid (HA) in JUVÉDERM® XC.


Please see additional Important Safety Information inside.

YOUR SKIN CHANGES OVER TIME

As you age, your skin loses elasticity and moisture. This normal aging process, plus environmental factors like sun exposure, cause different facial lines to form, including moderate to severe wrinkles and folds.



There are several different types of facial lines that form around your nose and mouth. These include the parentheses lines, vertical lip lines, corner lines, and marionette lines. As you age, these lines may become more noticeable.



“My inner impression of myself and my outer appearance didn't match.”



LEARN HOW TO DEFY LINES
AND GET NATURAL-LOOKING RESULTS



Actual patient.
Results may vary.

“Everyone will notice,
no one will know.”

— Shawn

HOW IT WORKS

JUVÉDERM® XC injectable gel:

- Is a nonsurgical treatment
- Contains a modified hyaluronic acid, derived from a naturally occurring sugar found in the human body
- Is gently injected by your healthcare professional under the skin using a fine needle
- Contains a common anesthetic called *lidocaine* to reduce discomfort

JUVÉDERM® XC is a nonsurgical filler your healthcare professional uses to instantly smooth away the deeper lines along the sides of your nose and mouth.

Get natural-looking results for up to 1 year with optimal treatment. Individual results may vary.

See the full gallery of JUVÉDERM® XC
before-and-after photos at Juvederm.com.

IMPORTANT SAFETY INFORMATION (continued)

What precautions should my doctor advise me about?

- The safety of JUVÉDERM® XC injectable gel for use during pregnancy, in women who are breastfeeding, or in patients with very thin skin in the cheek area has not been studied

IMPORTANT SAFETY INFORMATION (continued)

- The safety for use in patients under 18 years has not been studied
- The safety and effectiveness for treatment in areas other than facial wrinkles and folds (such as lips) have not been established

Please see additional Important Safety Information on following pages.

SEE THE DIFFERENCE WITH JUVÉDERM® XC

TALK TO YOUR HEALTHCARE PROFESSIONAL
TO SEE IF JUVÉDERM® XC IS RIGHT FOR YOU.

ABBEY



BEFORE

Individual results may vary.

AFTER

Unretouched photos of paid model taken before treatment and 2 weeks after treatment. A total of 3.2 mL of JUVÉDERM® XC was injected into the parentheses lines, corner lines, and marionette lines. These photos are not of a clinical trial subject.

LIZETTE



BEFORE

Individual results may vary.

AFTER

Unretouched photos of paid model taken before treatment and 2 weeks after treatment. A total of 2.0 mL of JUVÉDERM® XC was injected into the parentheses lines, corner lines, and marionette lines. These photos are not of a clinical trial subject.

JENNIFER



BEFORE

Individual results may vary.

AFTER

Unretouched photos of paid model taken before treatment and 2 weeks after treatment. A total of 3.0 mL of JUVÉDERM® XC was injected into the parentheses lines, corner lines, marionette lines, and vertical lip lines. These photos are not of a clinical trial subject.

LAURA



BEFORE

Individual results may vary.

AFTER

Unretouched photos of paid model taken before treatment and 2 weeks after treatment. A total of 4.2 mL of JUVÉDERM® XC was injected into the parentheses lines, corner lines, marionette lines, and vertical lip lines. These photos are not of a clinical trial subject.

IMPORTANT SAFETY INFORMATION (continued)

- The safety for use in patients with a history of excessive scarring or pigmentation disorders has not been studied and may result in additional scars or changes in pigmentation
- Tell your doctor if you are on therapy used to decrease the body's immune response (immunosuppressive therapy). Use may result in an increased risk of infection

IMPORTANT SAFETY INFORMATION (continued)

- Tell your doctor if you are using medications that can prolong bleeding, such as aspirin, ibuprofen, or other blood thinners; as with any injection, this may result in increased bruising or bleeding at the injection site

Please see additional Important Safety Information on following pages.

LONG-LASTING RESULTS UP TO 1 YEAR

The *first* and *only* injectable gel for wrinkles and folds that's FDA approved to **last up to 1 year** with optimal treatment.

LISA



BEFORE

AFTER 6 WEEKS

1 YEAR AFTER TREATMENT

Individual results may vary.

Unretouched photos of paid JUVÉDERM® XC injectable gel patient who received a 3-syringe treatment to correct parentheses lines. One syringe = 1.0 mL of JUVÉDERM® XC.

In clinical studies, optimal correction was achieved with 3.2 mL. Results lasted up to 1 year. These photos are not of a clinical trial subject.

EMILY



BEFORE

AFTER 6 WEEKS

1 YEAR AFTER TREATMENT

Individual results may vary.

Unretouched photos of paid JUVÉDERM® XC injectable gel patient who received a 3-syringe treatment to correct parentheses lines. One syringe = 1.0 mL of JUVÉDERM® XC.

In clinical studies, optimal correction was achieved with 3.2 mL. Results lasted up to 1 year. These photos are not of a clinical trial subject.

JUVÉDERM® XC is the # 1 selling filler in the US

It's your time to defy.

IMPORTANT SAFETY INFORMATION (continued)

- If laser treatment, chemical peel, or any other procedure based on active dermal response is considered after treatment, or if the product is administered before the skin has healed completely, there is a possible risk of an inflammatory reaction at the treatment site
- As with all skin injection procedures there is a risk of infection

Please see additional Important Safety Information on back page.

Visualize the difference for yourself with JUVÉDERM® XC



DOWNLOAD THE JUVÉDERM® XC TREATMENT VISUALIZER APP.*

The JUVÉDERM® XC Treatment Visualizer app is available for all Apple devices, as well as the *Android*™ smartphone.

*The JUVÉDERM® XC Treatment Visualizer app is for illustrative and demonstrative purposes only. The results displayed may not reflect actual results.

IMPORTANT SAFETY INFORMATION (continued)

What are possible side effects?

The most commonly reported side effects for JUVÉDERM® XC injectable gel were temporary injection-site redness, swelling, pain/tenderness, firmness, lumps/bumps, bruising, itching, and discoloration. They were predominantly mild or moderate in nature and lasted 7 days or less.

To report a problem with JUVÉDERM® XC, please call Allergan Product Surveillance at 1-800-624-4261.

For more information, please see the About Safety page at www.juvederm.com or call the Allergan Medical Information line at 1-800-433-8871.

JUVÉDERM® XC injectable gel is available by prescription only.

Ask your healthcare professional
if JUVÉDERM® XC is right for you.



©2014 Allergan, Inc. ® and ™ marks owned by Allergan, Inc.
JUVÉDERM® mark owned by Allergan Industrie, SAS.
Android is a trademark of Google Inc.
www.juvederm.com Re-order: APC37NC14 141249



Actual patient.
Results may vary.

DEFY GRAVITY.

A MORE YOUTHFUL PROFILE
STARTS WITH THE APPLES OF YOUR CHEEKS.

- **Instantly** adds volume lost to aging in the cheek area
- Creates **contour and a subtle lift**
- **Natural-looking results last up to 2 years** with optimal treatment

APPROVED USE

JUVÉDERM VOLUMA® XC injectable gel is for deep injection in the cheek area to correct age-related volume loss in adults over the age of 21.

IMPORTANT SAFETY INFORMATION

Are there any reasons why I should not receive JUVÉDERM VOLUMA® XC?

Do not use if you have a history of severe allergies or severe allergic reactions (anaphylaxis), or if you are allergic to the proteins (gram-positive bacterial proteins) used to make the hyaluronic acid (HA) in JUVÉDERM VOLUMA® XC.

Please see additional Important Safety Information inside.

WRINKLES AREN'T THE ONLY SIGN OF AGING

IS VOLUME LOSS IN THE CHEEKS MAKING YOU LOOK OLDER?

As you age, it's not just about lines and wrinkles. Your cheeks also lose volume and the skin may sag.

That's why a more youthful profile *starts* with the apples of your cheeks.



IMPORTANT SAFETY INFORMATION (continued)

What precautions should my doctor advise me about?

- The safety of JUVÉDERM VOLUMA® XC injectable gel for use during pregnancy, in women who are breastfeeding, or in patients with very thin skin in the cheek area has not been studied
- The safety for use in patients under 35 years or over 65 years has not been studied
- The safety and effectiveness for treatment in areas other than the cheek area have not been established



YOUTHFUL CHEEK:
Round, like an apple



AGING CHEEK:
Flat and sunken



V O L U M A[®] XC

IMPORTANT SAFETY INFORMATION (continued)

- The safety for use in patients with a history of excessive scarring or pigmentation disorders has not been studied and may result in additional scars or changes in pigmentation
- The long-term safety of repeat treatments has not been established

Please see additional Important Safety Information on the following pages.



DEFY GRAVITY

(SORRY, NEWTON. NOT EVERYTHING HAS TO FALL.)

DISCOVER THE **FIRST** AND **ONLY** FDA-APPROVED
INJECTABLE GEL FOR THE CHEEK AREA

- Adds volume lost to aging for adults over 21 years
- Creates **contour and a subtle lift instantly**
- Helps restore a **more youthful profile for up to 2 years** with optimal treatment
- Provides **natural-looking** results
- Is a **nonsurgical treatment**
- Contains a modified hyaluronic acid (HA) made from a naturally occurring complex sugar

Individual results may vary.




V O L U M A[®] XC

IMPORTANT SAFETY INFORMATION (continued)

- Patients who experience skin injury near the site of injection may be at a higher risk for side effects
- Tell your doctor if you are on therapy used to decrease the body's immune response (immunosuppressive therapy). Use may result in an increased risk of infection

HOW IT WORKS

JUVÉDERM VOLUMA® XC is injected under the skin using a fine needle. It can instantly add just enough volume to gently lift the cheek area.




To reduce discomfort during the procedure, JUVÉDERM VOLUMA® XC contains an anesthetic called *lidocaine*.

ASK YOUR HEALTHCARE PROFESSIONAL
ABOUT JUVÉDERM VOLUMA® XC TODAY.

IMPORTANT SAFETY INFORMATION (continued)

- Tell your doctor if you are using medications that can prolong bleeding, such as aspirin, ibuprofen, or other blood thinners; as with any injection, this may result in increased bruising or bleeding at the injection site
- Minimize strenuous exercise and exposure to extensive sun or heat within the first 24 hours following treatment

Please see additional Important Safety Information on the following pages.



“I look like
me, still,
but more
youthful.”

—Mara

Look up to 5 years younger
at 6 months after treatment.

As reported by people in the clinical study. Results may vary.



V O L U M A[®] XC

IMPORTANT SAFETY INFORMATION (continued)

What are possible side effects?

Side effects are moderate (uncomfortable) and generally last 2 to 4 weeks. The most common side effects include temporary reactions at the treatment site such as tenderness, swelling, firmness, lumps/bumps, bruising, pain, redness, discoloration, and itching.

To report a side effect, please call Allergan Product Surveillance at 1-877-345-5372.

For more information, please see the About Safety page at www.juvederm.com or call the Allergan Medical Information line at 1-800-766-0171.

JUVÉDERM VOLUMA[®] XC injectable gel is available by prescription only.

THE RESULTS: HIGHLY SATISFYING

“My cheeks look fuller in a subtle, natural way.”

—Mary Ann



BEFORE

AFTER



Individual results may vary.

Unretouched photos of paid model taken before treatment and 1 month after treatment. A total of 2.6 mL of JUVÉDERM VOLUMA® XC was injected into the cheek area.

The average volume of JUVÉDERM VOLUMA® XC injected during the clinical study was 6.8 mL to achieve optimal correction for the cheek area.

90% of people were more satisfied with their facial appearance at 6 months.
(n = 205)

As reported by people in the clinical study.

APPROVED USE

JUVÉDERM VOLUMA® XC injectable gel is for deep injection in the cheek area to correct age-related volume loss in adults over the age of 21.

IMPORTANT SAFETY INFORMATION

Are there any reasons why I should not receive JUVÉDERM VOLUMA® XC?

Do not use if you have a history of severe allergies or severe allergic reactions (anaphylaxis), or if you are allergic to the proteins (gram-positive bacterial proteins) used to make the hyaluronic acid (HA) in JUVÉDERM VOLUMA® XC.

Please see additional Important Safety Information on the inside panels.

THE RESULTS: NATURAL LOOK AND FEEL



“I see the
natural-looking
results I wanted.”

—Christine

BEFORE



AFTER



Individual results may vary.

Unretouched photos of paid model taken before treatment and 1 month after treatment. A total of 5.8 mL of JUVÉDERM VOLUMA® XC was injected into the cheek area.

IMPORTANT SAFETY INFORMATION (continued)

What precautions should my doctor advise me about?

- The safety of JUVÉDERM VOLUMA® XC injectable gel for use during pregnancy, in women who are breastfeeding, or in patients with very thin skin in the cheek area has not been studied
- The safety for use in patients under 35 years or over 65 years has not been studied
- The safety and effectiveness for treatment in areas other than the cheek area have not been established

FROM THE MAKERS OF THE #1 SELLING FAMILY OF FILLERS



Individual results may vary.

The average volume of JUVÉDERM VOLUMA® XC injected during the clinical study was 6.8 mL to achieve optimal correction for the cheek area.

 Juvéderm
V O L U M A[®] X C

IMPORTANT SAFETY INFORMATION (continued)

- The safety for use in patients with a history of excessive scarring or pigmentation disorders has not been studied and may result in additional scars or changes in pigmentation
- The long-term safety of repeat treatments has not been established

THE RESULTS: LONG LASTING



“I’ll grow old gracefully, but not passively.”

—Leslie

BEFORE



AFTER



Individual results may vary.

Unretouched photos of paid model taken before treatment and 1 month after treatment. A total of 4.0 mL of JUVÉDERM VOLUMA[®] XC was injected into the cheek area.


76% of people still experienced increased satisfaction **2 years** after treatment.
(n = 157)

As reported by people in the clinical study.

IMPORTANT SAFETY INFORMATION (continued)

- Patients who experience skin injury near the site of injection may be at a higher risk for side effects
- Tell your doctor if you are on therapy used to decrease the body's immune response (immunosuppressive therapy). Use may result in an increased risk of infection

LASTS UP TO 2 YEARS WITH OPTIMAL TREATMENT



Individual results may vary.

The average volume of JUVÉDERM VOLUMA® XC injected during the clinical study was 6.8 mL to achieve optimal correction for the cheek area.

SEE THE FULL GALLERY OF PHOTOS AT
JUVEDERM.COM

IMPORTANT SAFETY INFORMATION (continued)

- Tell your doctor if you are using medications that can prolong bleeding, such as aspirin, ibuprofen, or other blood thinners; as with any injection, this may result in increased bruising or bleeding at the injection site
- Minimize strenuous exercise and exposure to extensive sun or heat within the first 24 hours following treatment

Please see additional Important Safety Information on the back cover.

IT'S TIME TO DEFY GRAVITY

JUVÉDERM VOLUMA[®] XC instantly adds volume lost to aging in the cheek area. **Natural-looking** results **last up to 2 years** with optimal treatment.

 Juvéderm
V O L U M A[®] X C

IMPORTANT SAFETY INFORMATION (continued)

What are possible side effects?

Side effects are moderate (uncomfortable) and generally last 2 to 4 weeks. The most common side effects include temporary reactions at the treatment site such as tenderness, swelling, firmness, lumps/bumps, bruising, pain, redness, discoloration, and itching.

To report a side effect, please call Allergan Product Surveillance at 1-877-345-5372.

For more information, please see the About Safety page at www.juvederm.com or call the Allergan Medical Information line at 1-800-766-0171.

JUVÉDERM VOLUMA[®] XC injectable gel is available by prescription only.

ASK ABOUT
JUVÉDERM VOLUMA[®] XC TODAY.